

GROUP 2: WITHOUT PARTIAL VEIL, NOR RING, NON BULBOUS STIPE, STIPE EMERGING FROM A VOLVA LIKE A BAG OR A GLOVE FINGER.



Group 2A
White or gray cap, stipe emerging from a white or a gray volva.
2 species.



Text version





				<p>2,1 WHITE GRISETTE <i>AMANITA ALBIDICEPS</i></p>  
				<p>2,2 GRISETTE <i>AMANITA FUSCOZONATA</i></p>  

GROUP 2: WITHOUT PARTIAL VEIL, NOR RING, NON BULBOUS STIPE, STIPE EMERGING FROM A VOLVA LIKE A BAG OR A GLOVE FINGER.



Group 2A
White or gray cap, stipe emerging
from a white or a gray volva.
2 species.



<p>This medium-sized amanita is found mainly in poplar plantations and also in mixed forests, in summer and early autumn. It is distinguished from deadly white amanita by its non-bulbous stipe surrounded at the base by a sack-like volva. Its stipe don't wear a secondary veil, its bare cap (or adorned with some white velar remains) is ocher yellow in the center, the margin is long lined and the short gills are truncated.</p>	<p>2,1 WHITE GRISETTE <i>AMANITA ALBIDICEPS</i></p>  
<p>This name possibly includes about ten species or subspecies. The caps range from very pale gray to slate gray. In some varieties the volva is completely white, in others it is dark gray. The distinction between these different species can only be done by DNA testing.</p>	<p>2,2 GRISETTE <i>AMANITA FUSCOZONATA</i></p> 